



3.3

3.4

(mg) / (g) .
FVP
FVP
(P<0.05);
FVP
(P<0.05).

(MDA) :
MDA
MDA
(P<0.05);
MDA
(P<0.05).

Tab. 2 Spleen index of the mice in each group (n=10, x±s)

	3.55±0.74
	6.06±4.36
	4.65±1.19
FVP	5.63±4.14
FVP	4.86±1.75
FVP	4.01±0.91

(GSH-Px) :
GSH-Px
FVP
(P<0.05);
GSH-Px
(P<0.05).
(SOD) :
FVP
FVP
(P<0.05);
SOD
SOD
(P<0.05);
SOD
(P<0.05).

Tab. 3 Antioxidant indexes of the mice in each group (n=10, x±s, λ/(U·mg⁻¹))

	MDA	GSH-Px	SOD
	7.76±0.47	201.75±64.93	40.00±2.02
	18.01±0.58	154.16±14.80	25.22±3.58
	3.81±0.66	332.87±21.83	38.12±0.66
FVP	6.61±0.61	213.46±20.03	41.83±7.14
FVP	3.61±0.55	329.78±30.93	95.51±6.63
FVP	7.05±5.30	394.57±34.46	119.58±7.53

4

D-
[6] (MDA)
[4]
[5] (GSH-Px)
:6 (SOD)
D-
FVP [7]
(P<0.05), FVP FVP 0.05); MDA (P<
(P<0.05), (P<0.05), GSH-Px
SOD

FVP FVP SOD
 (P<0.05);
 SOD (P<0.05).
 D-

(*Flammulina velutipes*) extract added to beef and fish meats [J]. *Ajric Food Chem*, 2008, 1256 (21): 10032-10040.

[3] [D]. , 2009: 21-50.

[4] Wenjian Yang, Yong Fang, Jin Liang, *et al.* Optimization of ultrasonic extraction of *Flammulina velutipes* polysaccharides and evaluation of its acetylcholinesterase inhibitory activity[J]. *Food Research International*, 2011, 44: 1269-1275.

[5] [J]. , 2009(5): 72-73.

[6] [J]. (), 2002, 21 (50): 217.

[7] [M]. : , 2002: 50-149.

[8] [J]. , 2011, 29(4): 90-94.

[1] [J]. , 2008, 14(16): 2436-2438.
 [2] Bao H N, Ushio H, Ohshima T. Antioxidative activity and antidiscoloration efficacy of ergothioneine in mushroom

2014

	2 743	1		599	18
	2 357	2		552	19
	1 915	3	JOURNAL OF ZHEJIANG UNIVERSITY SCIENCE B	541	20
	1 490	4		529	21
	1 442	5		518	22
	1 319	6		516	23
	1 216	7		514	24
	1 202	8		476	25
	985	9		447	26
	922	10		429	27
	880	11		420	28
	829	12		407	29
	677	13		398	30
	665	14		386	31
	650	15	JOURNAL OF ZHEJIANG UNIVERSITY SCIENCE A	382	32
	645	16		
	614	17	62		